

MIT OpenCourseWare
<http://ocw.mit.edu>

18.02 Multivariable Calculus
Fall 2007

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

18.02 Practice Exam 2 A

Problem 1. (10 points: 5, 5)


Let $f(x, y) = xy - x^4$.

- Find the gradient of f at $P : (1, 1)$.
- Give an approximate formula telling how small changes Δx and Δy produce a small change Δw in the value of $w = f(x, y)$ at the point $(x, y) = (1, 1)$.

Problem 2. (20 points)

On the topographical map below, the level curves for the height function $h(x, y)$ are marked (in feet); adjacent level curves represent a difference of 100 feet in height. A scale is given.

- Estimate to the nearest .1 the value at the point P of the directional derivative $\left(\frac{dh}{ds}\right)_{\hat{u}}$, where \hat{u} is the unit vector in the direction of $\hat{i} + \hat{j}$.
- Mark on the map a point Q at which $h = 2200$, $\frac{\partial h}{\partial x} = 0$ and $\frac{\partial h}{\partial y} < 0$. Estimate to the nearest .1 the value of $\frac{\partial h}{\partial y}$ at Q .




Problem 3. (10 points)

Find the equation of the tangent plane to the surface $x^3y + z^2 = 3$ at the point $(-1, 1, 2)$.

Problem 4. (20 points: 5,5,5,5)

A rectangular box is placed in the first octant as shown, with one corner at the origin and the three adjacent faces in the coordinate planes. The opposite point $P : (x, y, z)$ is constrained to lie on the paraboloid $x^2 + y^2 + z = 1$. Which P gives the box of greatest volume?

- Show that the problem leads one to maximize $f(x, y) = xy - x^3y - xy^3$, and write down the equations for the critical points of f .
- Find a critical point of f which lies in the first quadrant ($x > 0, y > 0$).
- Determine the nature of this critical point by using the second derivative test.
- Find the maximum of f in the first quadrant (justify your answer).



Problem 5. (15 points)

In Problem 4 above, instead of substituting for z , one could also use Lagrange multipliers to maximize the volume $V = xyz$ with the same constraint $x^2 + y^2 + z = 1$.

- Write down the Lagrange multiplier equations for this problem.
- Solve the equations (still assuming $x > 0, y > 0$).

Problem 6. (10 points)

Let $w = f(u, v)$, where $u = xy$ and $v = x/y$. Using the chain rule, express $\frac{\partial w}{\partial x}$ and $\frac{\partial w}{\partial y}$ in terms of x, y, f_u and f_v .

Problem 7. (15 points)

Suppose that $x^2y + xz^2 = 5$, and let $w = x^3y$. Express $\left(\frac{\partial w}{\partial z}\right)_y$ as a function of x, y, z , and evaluate it numerically when $(x, y, z) = (1, 1, 2)$.